

W-algebras arising as chiral algebras of conformal field theory¹H. G. KAUSCH²*Theoretical Physics, Blackett Laboratory, Prince Consort Road,
Imperial College, London SW7 2BZ, U.K.*

It is argued that chiral algebras of conformal field theory possess a W-algebra structure. A survey of explicitly known W-algebras and their constructions is given.

Introduction

The presence of infinite dimensional *chiral* symmetries is central to the study of two dimensional conformal field theories. These symmetries, called *chiral algebras*, are formed by the purely holomorphic and purely anti-holomorphic fields, respectively. How the field content of a conformal field theory (CFT) is organised into representations of the chiral algebras is encoded in the fusion ring. The task of solving and classifying general conformal field theories can be split into two separate problems: to find chiral algebras of conformal field theory and to determine possible fusion rings of chiral algebras.

This programme has been most successful for the Virasoro minimal series [1], where the fields fall into finitely many representations of the Virasoro algebra. This has subsequently enabled the complete characterisation of the field content of such theories [2]. But considerable progress has also been made in constructing other chiral algebras of CFT. I present here a survey of these results.

Chiral algebras

The (left) chiral algebra of a two dimensional conformal field theory (CFT) is the symmetry algebra formed by all fields $\phi(z)$ which depend analytically on the coordinates. Locality implies that $[\phi(z), \psi(\zeta)] = 0$, for $z \neq \zeta$ and that all correlation functions are meromorphic in the arguments with poles only at coinciding arguments. From this one can derive the short distance behaviour of chiral fields which is given by the operator product expansion (OPE),

$$\phi(z)\psi(\zeta) = \sum_n (z - \zeta)^n \chi_n(\zeta), \quad (1)$$

where the $\chi_n(\zeta)$ are again local fields and only finitely many singular terms appear. Fields which transform homogeneously under projective transformations $\gamma: z \mapsto (az+b)/(cz+d)$ are

¹To appear in the proceedings of the XIX International Colloquium on "Group Theoretical Methods in Physics", Salamanca, Spain, June 29 – July 4, 1992

²Email: H.G.KAUSCH@V1.PH.IC.AC.UK

called quasi-primary (or non-derivative). They correspond to highest weights of the $su(1,1)$ algebra generated by the modes $L_0, L_{\pm 1}$ of the stress tensor. All other fields can be obtained by taking derivatives. We will thus restrict attention to quasi-primary fields and choose a basis $\{\chi^i\}$ with conformal weights $\{h_i\}$. Using $su(1,1)$ -covariance and the usual contour deformation argument we find from the OPE that the commutator of two quasi-primary fields has the form

$$[\chi_m^i, \chi_n^j] = \sum_k C_k^{ij} p_{h_k}^{h_i h_j}(m, n) \chi_{m+n}^k + D^{ij} \binom{m+h_i-1}{2h_i-1} \delta_{m+n}. \quad (2)$$

The contributions from a particular $su(1,1)$ representation are collected in the $su(1,1)$ -Clebsch-Gordon coefficients $p_{h_k}^{h_i h_j}(m, n)$ which are polynomials in m and n depending only on the conformal weights of the fields involved [3]. The $D^{ij} = \langle \chi^i | \chi^j \rangle$ define a metric on the space of quasi-primary fields which can be used to raise and lower indices. The field content of a particular chiral algebra is encoded in the structure constants D^{ij} and $C^{ijk} = \langle \chi^i | \chi^j(1) | \chi^k \rangle$. Locality of the chiral algebra is equivalent to the Jacobi identity for the commutator algebra which takes the form

$$\sum_{N=1}^{h_k+h_l-1} \left(\sum_{q,q'} C^{ijq} C^{klq'} D_{qq'} \right) p_{h_i}^{h_j h_q}(m, n+p) p_{h_q}^{h_k h_l}(n, p) + \text{cyclic} = 0. \quad (3)$$

The study of chiral algebras is thus equivalent to the study of the Lie algebra of quasi-primary fields. However, we still have to deal with an infinite number of quasi-primary fields.

In addition to the Lie bracket structure, which is determined solely by the *singular* part of the OPE, chiral algebras admit a second structure encoded in the *regular* part of the OPE, the so-called normal ordered products (NOPs). They are usually defined as the constant term of the OPE, $(\phi\psi)(z) = \oint_z \frac{d\zeta}{2\pi i} (\zeta - z)^{-1} \phi(\zeta) \psi(z)$. However, $(\phi\psi)(z)$ is not quasi-primary. We therefore define a new NOP, $\phi \diamond \psi$, by projecting $(\phi\psi)(z)$ onto quasi-primary fields, i.e. $\phi \diamond \psi$ is the new quasi-primary field entering the OPE of ϕ and ψ at the constant term. The $\diamond$ -product is commutative, but still non-associative, and differs from the usual NOP only by a finite number of correction terms which are known explicitly [3].

We call fields which can be obtained from the $\diamond$ -product *composite*. Fields which are orthogonal to all composite fields are called *simple*. The commutators of simple fields yields fields which are in general not simple but involve NOPs of simple fields. The algebraic structure given by commutators and NOPs of simple fields is called a *W-algebra*.

The simple fields generate the complete chiral algebra through NOPs and derivatives and all structure constants involving composite fields are determined by the couplings between simple fields. This means that the chiral algebra is completely determined by the W-algebra. Typically, a W-algebra has only a finite number of simple fields reducing the problem of constructing a chiral algebra to the *finite* problem of finding solutions to the associativity constraints (3).

One starts by specifying the weights of simple fields, $(h_1, \dots, h_r)$ and then attempts to construct the W-algebra $W_{(h_1, \dots, h_r)}$ as follows.

- Construct all composite fields of weight less than twice the maximum of the h_i . These are the only fields which can appear in the commutator of two simple fields.
- Express the structure constants for composite fields in terms of the structure constants for simple fields only.
- Impose the Jacobi identity for simple fields.

This yields a system of polynomial equations for the simple structure constants which one now has to solve. There are three possibilities:

- $W_{(h_1, \dots, h_r)}$ is consistent for generic values of c (*generic W-algebras*),
- $W_{(h_1, \dots, h_r)}$ is consistent for a finite set of c -values (*exceptional W-algebras*),
- $W_{(h_1, \dots, h_r)}$ is inconsistent for all values of c .

This approach is systematic enough to be amenable to algebraic computing. The results obtained so far [3–5] fall into the following classes.

Generic W-algebras

CURRENT ALGEBRAS: If the simple fields are all of weight one they form a current algebra. Their commutator algebra is a Kac-Moody algebra and the W-algebra is given by all derivatives and NOPs of the currents.

TODA THEORIES: W-algebras also appear as the chiral symmetry algebras of Toda field theory. The W-algebra fields are normal ordered differential polynomials of the Toda fields which commute with each of the exponential terms in the Toda Lagrangian [6]. The W-algebra Wg obtained from Toda field theory based on a Lie algebra g has rank g simple fields. Their weights are equal to the orders of the independent Casimir operators of g [7]. The following quantum W-algebras have been constructed so far:

Wg	weights	Wg	weights
WA_1	2	WG_2	2, 6
$W(A_1 \times U(1))$	2, 1	WA_3	2, 3, 4
$W(A_1 \times A_1)$	2, 2	WB_3	2, 4, 6
WA_2	2, 3	WC_3	2, 4, 6
WC_2	2, 4	$W(A_1^n)$	2^n

It has been realised recently that Toda field theories can be viewed as Hamiltonian reductions of constrained WZNW theories [8]. In this context it is possible to associate (at least classically) a W-algebra to every integral $sl(2)$ embedding into a simple Lie algebra. In this case the weights of the simple fields are one higher than the exponents of g with respect to the $sl(2)$ embedding. However, no quantum version of such a generalised algebra has been constructed as yet.

ORBIFOLD MODELS: Whenever a W-algebra has a (discrete) automorphism σ the subspace of fields invariant under σ forms again a W-algebra. An example of this is given by one of the algebras $W_{(2,4,6)}$, which turns out to be the bosonic projection of the super-Virasoro algebra [9].

Exceptional W-algebras

EXTENSIONS: If for a specific c -value a generic W-algebra possesses representations of integral weight with appropriate fusion rules then it is possible to enlarge the generic W-algebra by these representations to obtain an exceptional W-algebra. This is indicated by the appearance of a non-diagonal modular invariant [3]. The simplest example is given by the $(A_{p-1}, D_{q/2+1})$ modular invariant of the Virasoro algebra. Here, the field $\phi_{p-1,q-1}$ has weight $h = (p-2)(q-2)/4$ and fusion rule $\phi_{p-1,q-1} \times \phi_{p-1,q-1} = \mathbf{1}$. Whenever p or q is even, this yields a W-algebra $W_{(2,h)}$. Exceptional W-algebras coming from such an extension of the Virasoro algebra are listed below. Extensions of other generic W-algebras have not yet been constructed.

weights	c -values	weights	c -values
2, 2	$-\frac{44}{5}, -\frac{39}{10}, -\frac{494}{35}$	2, 7	$-\frac{25}{2}$
2, 3	$-\frac{114}{7}, -2, \frac{4}{5}$	2, 8	$-\frac{944}{17}, \frac{21}{22}, -\frac{224}{65}$
2, 4	$-\frac{11}{14}$	2, 3, 3, 3	-2
2, 5	$-\frac{350}{11}, -7, \frac{6}{7}$	2, 5, 5, 5	-7
2, 6	$-\frac{516}{13}, -\frac{306}{55}$	2, 7, 7, 7	$-\frac{25}{2}$

DECOUPLING: It can happen that at specific values of c some of the simple fields of a generic W-algebra decouple and one is left with an exceptional W-algebra. The simplest example is the Ising-model $c = \frac{1}{2}$ which is in the WE_8 minimal series. In this case all simple fields apart from the stress tensor decouple. Known exceptional W-algebras arising from this decoupling procedure are:

Wg	weights	$\longrightarrow$	weights	c -values
WE_6	2, 5, 6, 8, 9, 12	$\longrightarrow$	2, 5	$-\frac{350}{11}, -7, \frac{6}{7}$
WE_8	2, 8, 12, 14, 18, 20, 24, 30	$\longrightarrow$	2, 8	$-\frac{944}{17}$
WD_4	2, 4, 4, 6	$\longrightarrow$	2, 4, 4	$1, -\frac{656}{11}$

These results support the conjecture that all generic W-algebras are all given by Hamiltonian reduction of constrained WZW models and their orbifolds. The picture for exceptional W-algebras is much more complicated. While the majority of cases fit the two procedures of extending and decoupling there are a large number of cases which do not fall into these patterns [3, 10].

References

- [1] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, Nucl. Phys. B241 (1984) 333.
- [2] A. Cappelli, C. Itzykson and J.-B. Zuber, Commun. Math. Phys. 113 (1987) 1.
- [3] R. Blumenhagen, M. Flohr, A. Kliem, W. Nahm, A. Recknagel and R. Varnhagen, Nucl. Phys. B361 (1991) 255.
- [4] H. G. Kausch and G. M. T. Watts, Nucl. Phys. B354 (1991) 740.
- [5] H. G. Kausch, Phys. Lett. 259B (1991) 448.
- [6] H. G. Kausch and G. M. T. Watts, Int. J. Mod. Phys. A7 No. 17 (1992) 4175.
- [7] B. L. Feigin and E. V. Frenkel, Int. Math. Research Notices 1991 No. 6 75.
- [8] L. Fehér, L. O’Raifeartaigh, L. Ruelle, I. Tsutsui and A. Wipf, On the general structure of Hamiltonian reductions of the WZNW theory, DIAS-STP-91-29.
- [9] P. Bouwknegt, Extended conformal algebras from Kac-Moody algebras, *in*: Infinite Dimensional Lie Algebras and Lie Groups, ed. V. G. Kac, World Scientific, 1988, CIRM-Luminy July meeting, Marseille 1988.
- [10] W. Eholzer, M. Flohr, A. Honecker, R. Hübel, W. Nahm and R. Varnhagen, Representations of W-algebras with two generators and new rational models, BONN-HE-92-02, to appear in Nucl. Phys. B.